

**American College of Radiology
ACR Appropriateness Criteria®**

Clinical Condition: **Right Lower Quadrant Pain**

Variant 1: **Fever, leukocytosis, and classic presentation clinically for appendicitis in adults.**

Radiologic Procedure	Rating	Comments	RRL*
CT abdomen and pelvis with contrast	8		High
US abdomen RLQ graded compression	6		None
CT abdomen and pelvis without contrast	6		High
X-ray chest	5		Min
US pelvis transabdominal and transvaginal	5		None
X-ray abdomen supine and upright	5		Low
X-ray colon barium enema double-contrast	4		Med
X-ray colon barium enema single-contrast	4		Med
MRI abdomen and pelvis	4		None
X-ray small bowel series with barium	3		Low
NUC gallium scan abdomen	3		High
NUC WBC scan abdomen pelvis	3		Med
X-ray small bowel enteroclysis	2		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Right Lower Quadrant Pain****Variant 2:****Fever, leukocytosis; possible appendicitis, atypical presentation, adults and adolescents.**

Radiologic Procedure	Rating	Comments	RRL*
CT abdomen and pelvis with contrast	8		High
X-ray abdomen supine and upright	6		Low
US abdomen RLQ graded compression	6		None
US pelvis transabdominal and transvaginal	6		None
CT abdomen and pelvis without contrast	6		High
X-ray chest	5		Min
X-ray colon barium enema double-contrast	5		Med
X-ray colon barium enema single-contrast	5		Med
MRI abdomen and pelvis	5		None
X-ray small bowel series with barium	4		Low
NUC gallium scan abdomen	3		High
NUC WBC scan abdomen pelvis	3		Med
X-ray small bowel enteroclysis	2		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Right Lower Quadrant Pain****Variant 3:****Fever, leukocytosis, pregnant woman.**

Radiologic Procedure	Rating	Comments	RRL*
US abdomen RLQ graded compression	8		None
MRI abdomen and pelvis	7		None
US pelvis transabdominal and transvaginal	6		None
CT abdomen and pelvis with contrast	6		High
CT abdomen and pelvis without contrast	5		High
X-ray chest	4		Min
X-ray abdomen supine and upright	2		Low
X-ray colon barium enema double-contrast	2		Med
X-ray small bowel enteroclysis	2		Med
X-ray colon barium enema single-contrast	2		Med
NUC WBC scan abdomen pelvis	2		Med
X-ray small bowel series with barium	2		Low
NUC gallium scan abdomen	2		High
Rating Scale: 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

Clinical Condition:**Right Lower Quadrant Pain****Variant 4:****Fever, leukocytosis, possible appendicitis, atypical presentation in children (less than 14 years of age).**

Radiologic Procedure	Rating	Comments	RRL*
US abdomen RLQ graded compression	8		None
CT abdomen and pelvis with contrast	7	May be useful following negative US.	High
X-ray abdomen supine and upright	6		Low
US pelvis transabdominal and transvaginal	5		None
X-ray chest	5		Min
CT abdomen and pelvis without contrast	5		High
MRI abdomen and pelvis	5		None
X-ray colon barium enema single-contrast	3		Med
X-ray colon barium enema double-contrast	3		Med
X-ray small bowel series with barium	3		Low
NUC gallium scan abdomen	2		High
X-ray small bowel enteroclysis	2		Med
NUC WBC scan abdomen pelvis	2		Med
<u>Rating Scale:</u> 1=Least appropriate, 9=Most appropriate			*Relative Radiation Level

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

RIGHT LOWER QUADRANT PAIN

Expert Panel on Gastrointestinal Imaging: Robert L. Bree, MD, MHSA¹; C. Craige Blackmore, MD, MPH²; W. Dennis Foley, MD³; Spencer B. Gay, MD⁴; Seth N. Glick, MD⁵; Jay P. Heiken, MD⁶; James E. Huprich, MD⁷; Marc S. Levine, MD⁸; Pablo R. Ros, MD, MPH⁹; Max Paul Rosen, MD, MPH¹⁰; William P. Shuman, MD¹¹; Frederick L. Greene, MD¹²; Don C. Rockey, MD.¹³

Summary of Literature Review

Few comparative imaging studies evaluating right lower quadrant pain are available. Most imaging reports center on disease processes, such as appendicitis. Because appendicitis is the most common cause of right lower quadrant pain, the focus of this narrative is on appendicitis and the accuracy of imaging procedures in diagnosing appendicitis, although consideration of other diseases is, of course, included.

Acute appendicitis is the most common acute abdominal disorder that requires surgery. In most patients with acute appendicitis, imaging may not be necessary, because the clinical presentation is sufficiently diagnostic to allow surgery. To date, however, no prediction rules for identifying subjects with appendicitis have been validated. In the published studies for imaging in appendicitis, the selection criteria for imaging are not often stated, but in most investigations, subjects with definitive clinical exam findings of appendicitis undergo operation without imaging. In the reported imaging studies, an average of 45%-50% of imaged subjects had appendicitis, and 36% had nonspecific abdominal pain. Data on the overall effect of imaging on surgical treatment of appendicitis and patient outcome remain contradictory.

Plain film diagnosis is of limited value evaluating acute appendicitis, except in occasional circumstances when an appendicolith or other ancillary findings are identified. Although barium enema has been used historically to diagnose appendicitis, it depends on the negative finding of nonvisualization of the appendix and may be quite uncomfortable in patients with acute appendicitis.

Nonetheless, barium small-bowel follow-through or barium enema may be useful for other causes of right lower quadrant pain, including suspected small bowel obstruction, infectious ileitis, and inflammatory bowel disease. Finally, use of MRI for appendicitis has been reported in a few small case series, including in pregnant women.

Computed tomography (CT) is the most accurate study for evaluating patients without a clear clinical diagnosis of acute appendicitis. In a meta-analysis of prospective studies of the accuracy of CT and ultrasonography in adolescents and adults, CT demonstrated superior sensitivity (0.94, 95% CI: 0.91 to 0.95) and specificity (0.95, 95% CI: 0.93 to 0.96) versus ultrasound (US) (sensitivity 0.86, 95% CI: 0.83 to 0.88; specificity 0.81, 95% CI: 0.78 to 0.84). This analysis was based on 12 studies of CT and 14 studies of US identified through December 2003, and included four studies that directly compared both modalities. The results of investigations of CT showed consistent results across all studies and institutions, while US investigations demonstrated heterogeneity, suggesting greater dependence on operator skill.

Another controversy is whether or not to use intravenous contrast in the CT evaluation of appendicitis. High accuracy has been reported for both techniques, and direct comparisons are lacking. However, the majority of the available evidence is on CT with intravenous contrast. Institutional experience may be the best determinant appropriateness of intravenous contrast. Both CT and US may be effective in detecting causes of pain unrelated to appendicitis. CT has been reported to show a non-appendicitis cause of abdominal pain in 20% of subjects, versus 15% for US. The range of diseases studied includes inflammatory bowel disease, infectious bowel disease, small bowel obstruction, acute gynecological conditions and others.

CT appears superior to sonography in evaluating patients with periappendiceal abscess, especially when the abscesses become large. CT can be used to choose among different therapeutic options, including antibiotic treatment (with small abscesses), percutaneous drainage (with one to three well-defined medium-sized abscesses), and surgery (with extensive abnormality not amenable to percutaneous drainage).

CT and US have been less well evaluated in children than in adults. Many large prospective studies include subjects of all ages, despite the potential differences in imaging accuracy between children and adults due to smaller body size and less body fat in children. This makes it difficult

¹Review Author and Panel Chair, Radia Medical Imaging, Everett, Wash; ²Research Author, University of Washington, Seattle, Wash; ³Froedtert Hospital East, Milwaukee, Wis; ⁴University of Virginia Health Science Center, Charlottesville, Va; ⁵Presbyterian Medical Center, Philadelphia, Pa; ⁶Mallinckrodt Institute of Radiology, St. Louis, Mo; ⁷Mayo Clinic, Rochester, Minn; ⁸Hospital of the University of Pennsylvania, Philadelphia, Pa; ⁹Brigham & Women's Hospital, Boston, Mass; ¹⁰Beth Israel Hospital, Boston, Mass; ¹¹University of Washington, Seattle, Wash; ¹²Carolinas Medical Center, Charlotte, NC, American College of Surgeons; ¹³University of Texas, Southwest Medical Center, Dallas, Texas, American Gastroenterological Association.

Reprint requests to: Department of Quality & Safety, American College of Radiology, 1891 Preston White Drive, Reston, VA 20191-4397.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

to determine the accuracy of imaging in different subgroups. Further, the increased radiosensitivity of children makes the use of ionizing radiation of more concern for them. A systematic literature review through July 2004 revealed eight prospective evaluations of US for appendicitis in children. The pooled sensitivity of graded compression US was 91% (95% CI: 89%-93%), and the specificity was 97% (95% CI: 95%-99%). Only a single prospective study of CT in children was identified, reporting a sensitivity of 95%, and specificity of 98%. There is also a small literature on use of US as an initial imaging study, followed by CT for equivocal cases. Such combined protocols demonstrate sensitivity of 95% (95% CI: 83%-100%), and specificity 93% (95% CI: 87%-97%). These results suggest that although CT is more accurate, US may also be appropriate in experienced hands, particularly if equivocal results are followed up by CT.

Nuclear medicine imaging with WBC scans has also been reported for evaluating right lower quadrant pain.

Evaluation of the accuracy of imaging in pregnant women has received little attention in the literature. In general, ionizing radiation from CT should be avoided during pregnancy, and US is clearly a safer imaging option. However, with the absence of evidence, no specific recommendation can be made.

References

1. Addiss DG, Shaffer N, Fowler BS, Tauxe RV. The epidemiology of appendicitis and appendectomy in the United States. *Am J Epidemiol* 1990; 132(5):910-925.
2. Wagner JM, McKinney WP, Carpenter JL. Does this patient have appendicitis? *JAMA* 1996; 276(19):1589-1594.
3. Hallan S, Asberg A. The accuracy of C-reactive protein in diagnosing acute appendicitis--a meta-analysis. *Scand J Clin Lab Invest* 1997; 57(5):373-380.
4. Terasawa T, Blackmore CC, Bent S, Kohlwes RJ. Systematic review: computed tomography and ultrasonography to detect acute appendicitis in adults and adolescents. *Ann Int Med* 2004; 141(7):537-546.
5. Flum DR, Morris A, Koepsell T, Dellinger EP. Has misdiagnosis of appendicitis decreased over time? A population-based analysis. *JAMA* 2001; 286(14):1748-1753.
6. Garcia Pena BM, Mandl KD, Kraus SJ, et al. Ultrasonography and limited computed tomography in the diagnosis and management of appendicitis in children. *JAMA* 1999; 282(11):1041-1046.
7. Applegate KE, Sivit CJ, Salvator AE, et al. Effect of cross-sectional imaging on negative appendectomy and perforation rates in children. *Radiology* 2001; 220(1): 103-107.
8. Partrick DA, Janik JE, Janik JS, et al. Increased CT scan utilization does not improve the diagnostic accuracy of appendicitis in children. *J Pediatr Surg* 2003; 38(5):659-662.
9. Bendeck SE, Nino-Murcia M, Berry GJ, Jeffrey RB Jr. Imaging for suspected appendicitis: negative appendectomy and perforation rates. *Radiology* 2002; 225(1):131-136.
10. Rao PM, Rhea JT, Novelline RA, et al. Effect of computed tomography of the appendix on treatment of patients and use of hospital resources. *N Engl J Med* 1998; 338(3):141-146.
11. Balthazar EJ, Birnbaum BA, Yee J, et al. Acute appendicitis: CT and US correlation in 100 patients. *Radiology* 1994; 190(1):31-35.

12. Horton MD, Counter SF, Florence MG, Hart MJ. A prospective trial of computed tomography and ultrasonography for diagnosing appendicitis in the atypical patient. *Am J Surg* 2000; 179(5):379-381.
13. Kan JH, Fines BP, Funaki B. Conventional and hydrocolonic US of the appendix with CT correlation performed by on-call radiology residents. *Acad Radiol* 2001; 8(12):1208-1214.
14. Wise SW, Labuski MR, Kasales CJ, et al. Comparative assessment of CT and sonographic techniques for appendiceal imaging. *AJR* 2001; 176(4):933-941.
15. Lane MJ, Katz DS, Ross BA, et al. Unenhanced helical CT for suspected acute appendicitis. *AJR* 1997; 168(2):405-409.
16. Walker S, Haun W, Clark J, et al. The value of limited computed tomography with rectal contrast in the diagnosis of acute appendicitis. *Am J Surg* 2000; 180(6):450-454.
17. Wijetunga R, Tan BS, Rouse JC, et al. Diagnostic accuracy of focused appendiceal CT in clinically equivocal cases of acute appendicitis. *Radiology* 2001; 221(3):747-753.
18. Adams DH, Fine C, Brooks DC. High-resolution real-time ultrasonography. A new tool in the diagnosis of acute appendicitis. *Am J Surg* 1988; 155(1):93-97.
19. Bilbey JH, Gibney RG, Cooperberg PL. Ultrasonography in acute appendicitis. *Can Assoc Radiol J* 1989; 40(1):22-24.
20. Chen SC, Chen KM, Wang SM, Chang KJ. Abdominal sonography screening of clinically diagnosed or suspected appendicitis before surgery. *World J Surg* 1998; 22(5):449-452.
21. Incesu L, Coskun A, Selcuk MB, et al. Acute appendicitis: MR imaging and sonographic correlation. *AJR* 1997; 168(3):669-674.
22. Kang WM, Lee CH, Chou YH, et al. A clinical evaluation of ultrasonography in the diagnosis of acute appendicitis. *Surgery* 1989; 105(2 Pt 1):154-159.
23. Vermeulen B, Morabia A, Unger PF, et al. Acute appendicitis: influence of early pain relief on the accuracy of clinical and US findings in the decision to operate—a randomized trial. *Radiology* 1999; 210(3):639-643.
24. Baldisserotto, M, Marchiori E. Accuracy of noncompressive sonography of children with appendicitis according to the potential positions of the appendix. *AJR* 2000; 175(5):1387-1392.
25. Nunez D Jr, Yrizarry JM, Casillas VJ, et al. Percutaneous management of appendiceal abscesses. *Semin Ultrasound CT MR* 1989; 10(4):348-351.
26. Kuligowska E, Keller E, Ferrucci JT. Treatment of pelvic abscesses: value of one-step sonography guided transrectal needle aspiration and lavage. *AJR* 1995; 164(1):201-206.
27. Jeffrey RB Jr, Tolentino CS, Federle MP, Laing FC. Percutaneous drainage of periappendiceal abscesses: review of 20 patients. *AJR* 1987; 149(1):59-62.
28. Foley CR, Latimer RG, Rimkus DS. Detection of acute appendicitis by technetium 99 HMPAO scanning. *Am Surg* 1992; 58(12):761-765.
29. Puylaert JB. Acute appendicitis: US evaluation using graded compression. *Radiology* 1986; 158(2):355-360.
30. Puylaert JB, Rutgers PH, Lalisang RI, et al. A prospective study of ultrasonography in the diagnosis of appendicitis. *N Eng J Med* 1987; 317(11):666-669.
31. Schwerk WB, Wichtrup B, Rothmund M, Ruschoff J. Ultrasonography in the diagnosis of acute appendicitis: a prospective study. *Gastroenterology* 1989; 97(3):630-639.
32. Skaane P, Amland PF, Nordshus T, Solheim K. Ultrasonography in patients with suspected acute appendicitis: a prospective study. *Br J Radiol* 1990; 63(754):787-793.
33. Rao PM, Rhea JT, Rattner DW, et al. Introduction of appendiceal CT: impact on negative appendectomy and appendiceal perforation rates. *Ann Surg* 1999; 229(3):344-349.
34. Blackmore CC, Avey GD. Imaging in acute abdominal pain. In Medina LS, Blackmore CC, eds. *Evidence based imaging: Optimizing imaging for patient care*. New York, NY: Springer Verlag, 2006.
35. Lessin MS, Chan M, Catalozzi M, et al. Selective use of ultrasonography for acute appendicitis in children. *Am J Surg*

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.

- 1999; 177(3):193-196.
36. Hahn HB, Hoepner FU, Kalle T, et al. Sonography of acute appendicitis in children: 7 years experience. *Pediatr Radiol* 1998; 28(3): 147-151.
 37. Vignault F, Filiatrault D, Brandt ML, et al. Acute appendicitis in children: evaluation with US. *Radiology* 1990; 176(2):501-504.
 38. Quillin SP, Siegel MJ. Appendicitis: efficacy of color Doppler sonography. *Radiology* 1994; 191(2):557-560.
 39. Schulte B, Beyer D, Kaiser C, et al. Ultrasonography in suspected acute appendicitis in childhood-report of 1285 cases. *Eur J Ultrasound* 1998; 8(3):177-182.
 40. Kaiser S, Frenckner B, Jorulf HK. Suspected appendicitis in children: US and CT—a prospective randomized study. *Radiology* 2002; 223(3):633-638.
 41. Teo EL, Tan KP, Lam SL, et al. Ultrasonography and computed tomography in a clinical algorithm for the evaluation of suspected acute appendicitis in children. *Singapore Med J* 2000; 41(8):387-392.
 42. Lowe LH, Draud KS, Hernanz-Schulman M, et al. Nonenhanced limited CT in children suspected of having appendicitis: prospective comparison of attending and resident interpretations. *Radiology* 2001; 221(3):755-759.
 43. Lim HK, Bae SH, Seo GS. Diagnosis of acute appendicitis in pregnant women: value of sonography. *AJR* 1992; 159(3):539-542.

An ACR Committee on Appropriateness Criteria and its expert panels have developed criteria for determining appropriate imaging examinations for diagnosis and treatment of specified medical condition(s). These criteria are intended to guide radiologists, radiation oncologists and referring physicians in making decisions regarding radiologic imaging and treatment. Generally, the complexity and severity of a patient's clinical condition should dictate the selection of appropriate imaging procedures or treatments. Only those exams generally used for evaluation of the patient's condition are ranked. Other imaging studies necessary to evaluate other co-existent diseases or other medical consequences of this condition are not considered in this document. The availability of equipment or personnel may influence the selection of appropriate imaging procedures or treatments. Imaging techniques classified as investigational by the FDA have not been considered in developing these criteria; however, study of new equipment and applications should be encouraged. The ultimate decision regarding the appropriateness of any specific radiologic examination or treatment must be made by the referring physician and radiologist in light of all the circumstances presented in an individual examination.